KNEEbraska

Matthew R. Byington, DO Orthopaedic Surgeon—Board Certified

Sports Medicine & Arthroscopic Surgery // Knee & Shoulder Reconstruction
Prairie Orthopaedic & Plastic Surgery 4130 Pioneer Woods Dr, Suite 1, Lincoln, NE 68506

Phone: (402) 489-4700 Fax: (402) 489-5220

www.KNEEbraska.com // www.drmattbyington.com // www.prairie-ortho.com

LATERAL EPICONDYLITIS POST-OPERATIVE REHABILITATION PROTOCOL

	ROM	IMMOBILIZER	EXERCISE
PHASE I	PROM as tolerated	Worn for first 7-10	Gentle
(0-6 WEEKS)		days post-op	hand/wrist/elbow/shoulder
			strength work—no wrist
			extension
PHASE II	Increase ROM to	NONE	Advance exercise as
(6-8 WEEKS)	full, begin active wrist extension		above to resistive, maintain flexibility/ROM,
	Wilst Catolision		may begin gentle AROM
			wrist extension exercises
PHASE III	Full and pain-free	NONE	Advance exercise as
(8-10 WEEKS)			above, gradual
			progression to full activity

^{*}NO AROM wrist extension x 6 weeks post-operative