

KNEEbraska  
 Matthew R. Byington, DO  
 Orthopaedic Surgeon—Board Certified  
 Sports Medicine & Arthroscopic Surgery // Knee & Shoulder Reconstruction  
 Prairie Orthopaedic & Plastic Surgery 4130 Pioneer Woods Dr, Suite 1, Lincoln, NE 68506  
 Phone: (402) 489-4700 Fax: (402) 489-5220  
[www.KNEEbraska.com](http://www.KNEEbraska.com) // [www.drmmattbyington.com](http://www.drmmattbyington.com) // [www.prairie-ortho.com](http://www.prairie-ortho.com)

## REVERSE TOTAL SHOULDER REPLACEMENT REHABILITATION PROTOCOL

	<b>ROM</b>	<b>IMMOBILIZER</b>	<b>EXERCISES</b>
<b>PHASE I</b> (0-4 weeks)	Limit 45° passive ER to protect Subscapularis repair  Flexion , Extension progress as tolerated	<b>0-2 weeks:</b> Worn at all times day and night; off for gentle exercise only  <b>2-4 weeks:</b> Worn daytime only	<b>0-2 weeks:</b> Grip strengthening, pendulum exercises, elbow/wrist/ hand ROM at home  <b>2-4 weeks:</b> Begin cuff, deltoid isometrics; Limit ER to passive 45° and no active IR or extension until 6 weeks if Subscapularis repair was done
<b>PHASE II</b> (4-12 weeks)	Increase as tolerated  Begin active/active-assisted IR and extension as tolerated after 6 weeks if Subscapularis repair was done	NONE	<b>4-8 weeks:</b> Begin light resisted ER, forward flexion and abduction  Focus on Anterior Deltoid, Teres  <b>8-12 weeks:</b> Begin resisted IR, extension and scapular retraction
<b>PHASE III</b> (12-24 weeks)	Progress to full without discomfort	NONE	Advance strength training as tolerated; closed chain scapular rehab and functional rotator cuff strengthening; focus on anterior Deltoid and Teres mm. Emphasis scapular stabilization